



Title	Gut and oral microbiota in autoimmune liver disease
Author(s)	Abe, Kazumichi; Fujita, Masashi; Hayashi, Manabu; Okai, Ken; Takahashi, Atsushi; Ohira, Hiromasa
Citation	Fukushima Journal of Medical Science. 65(3): 71-75
Issue Date	2019
URL	http://ir.fmu.ac.jp/dspace/handle/123456789/1053
Rights	© 2019 The Fukushima Society of Medical Science. This article is licensed under a Creative Commons [Attribution-NonCommercial-ShareAlike 4.0 International] license.
DOI	10.5387/fms.2019-21
Text Version	publisher

[Review]

Gut and oral microbiota in autoimmune liver disease

Kazumichi Abe, Masashi Fujita, Manabu Hayashi, Ken Okai,
Atsushi Takahashi and Hiromasa Ohira

Department of Gastroenterology, Fukushima Medical University School of Medicine

(Received August 27, 2019, accepted September 2, 2019)

Abstract

The gut microbiota plays a key role in the development of chronic inflammatory liver disease. The gut-liver axis involves inflammatory cells, cytokines, and other molecules that cause liver deterioration. Dysbiosis is important in understanding several liver diseases, especially in relation to the development of autoimmune liver disease. The aim of this review is to provide a current overview of alterations in the gut and oral microbiota associated with autoimmune liver diseases.

Key words : gut microbiota, oral microbiota, autoimmune hepatitis, primary biliary cholangitis

Introduction

Primary biliary cholangitis (PBC) and autoimmune hepatitis (AIH) are classically viewed as distinct autoimmune liver diseases. Autoimmune liver diseases are thought to be triggered by environmental factors in genetically susceptible individuals. Genome-wide association and murine model studies have expanded our knowledge of autoimmune liver diseases, however, the pathogenesis of disease remains obscure. Recently culture-independent techniques have revolutionized knowledge of the gut and oral microbiota. This review provides an overview of gut and oral microbiota in autoimmune liver diseases.

Gut-Liver axis

The gut and liver are closely related. The liver is significantly affected by the gut and its contents, as 70% of the blood supplied to the liver is collected from intestinal circulation through the portal venous system. Both the gut and liver play a pivotal role in the absorption and metabolism of various nutrients and drugs. Abnormal bile acid homeostasis may lead to diarrhea and bacterial overgrowth¹⁾. Bacterial overgrowth, increased intestinal permeability, failure to inactivate endotoxins, and ac-

tivated innate immunity all contribute to the pathological state of bacterial translocation.

Gut and oral microbiota

The gut microbiota collectively refers to the 100 trillion bacteria, with an estimated mass of 1-2 kg, that inhabit the human gastrointestinal tract. This very diverse ecosystem comprises over 2,000 distinct species and has a collective genome containing 150-fold more genes than the human genome²⁾. The metabolic activity of the gut microbiota provides benefits to human health by supplying essential nutrients and maximizing the efficiency of energy harvest from ingested food. However, the microbiota also contains numerous potential opportunistic pathogens. Moreover, gut microbial products activate Toll-like receptors (TLRs) and induce inflammation that defines disease. Maintaining the homeostasis of the gut microbiota has thus necessitated the development of a specialized mucosal immune system³⁾. Characterization of the microbiota has been performed by culture and biochemical typing, which have been gold standards for the identification of bacterial species. Recently, however, advances in culture-independent techniques have revolutionized our knowledge about the gut microbiota. These techniques, based on sequence diver-

Corresponding author : Kazumichi Abe, MD E-mail : k-abe@fmu.ac.jp
<https://www.jstage.jst.go.jp/browse/fms> <http://www.fmu.ac.jp/home/lib/F-igaku/>

gences in the small subunit ribosomal ribonucleic acid (16S rRNA), allow the demonstration of gut microbial diversity, providing qualitative and quantitative information on bacterial species as well as alterations in the gut microbiota in healthy and diseased states⁴.

The oral cavity is a large reservoir of bacteria of more than 700 species or phylotypes and is profoundly relevant to host health and disease⁵⁻⁷. The role of the oral and gut microbiota in the pathogenesis of immune-related diseases has been highlighted in autoimmune diseases, such as autoimmune encephalomyelitis, rheumatoid arthritis, and inflammatory bowel disease⁸⁻¹⁴. A previous report revealed evidence of pervasive changes in the immune system-microbiota interface in the saliva of patients with cirrhosis that were similar to those found in stool¹⁵.

Gut and oral microbiota in AIH

AIH manifests as chronic liver inflammation of an unknown cause. It generally affects young to middle-aged females and is associated with the presence of autoantibodies and hypergammaglobulinemia¹⁶. However, the pathogenesis of this disease remains obscure. It is increasingly recog-

nized that the composition of the gut microbiota plays a critical role in influencing the predisposition to AIH¹⁷⁻²¹ (Table 1). In addition, an increase in the abundance of *Veillonella dispar* was associated with disease severity, and the metagenomic profile suggests that this bacterium may be involved in lipopolysaccharide (LPS) biosynthesis and amino acid metabolism²⁰. Lin R *et al.* reported that increased intestinal permeability and bacterial translocation occurred in AIH, which associated with disease severity¹⁷. However, direct evaluation of the oral microbiome has not been performed in AIH. Our data indicated a significant increase in the abundance of the genus *Veillonella* in the oral microbiota of patients with AIH, and the abundance of the genus *Veillonella* was positively correlated with the levels of proinflammatory cytokines, such as IL-1 β , IL-6, IL-8, and IL-12p70 in the saliva of patients with AIH²¹ (Figure 1a). However, it is unknown whether the inflammatory state in the oral cavity of AIH patients is the cause or consequence of imbalances in the salivary microbiota and whether the oral cavity or the gut immune response exerts greater influence on the observed dysbiosis of the oral microbiota. In our study, changes in the gut microbiota composition in AIH were characterized by an increase in abundance of the order *Lactobacillales* and

Table 1. Changes in the gut and oral microbiota associated with autoimmune liver diseases

Gut microbiota			
Disease	Comparison	Genus (or Order)	Author/year
AIH (n=24)	Healthy vs AIH	<i>Lactobacillus</i> ↓ <i>Bifidobacterium</i> ↓	Lin/2015 ¹⁷
AIH (n=17)	Healthy vs AIH	<i>Lactobacillales</i> ↑ <i>Clostridium</i> subcluster XIVa ↓	Abe/2018 ²¹
AIH (n=99)	Healthy vs AIH	<i>Veillonella</i> ↑, <i>Lactobacillus</i> ↑, <i>Streptococcus</i> ↑, <i>Klebsiella</i> ↑, <i>Oscillospira</i> ↓, <i>Parabacteroides</i> ↓, <i>Coprococcus</i> ↓	Wei/2019 ²⁰
PBC (n=79)	Healthy vs PBC	<i>Enterobacteriaceae</i> ↑ <i>Klebsiella</i> ↑, <i>Haemophilus</i> ↑ <i>Veillonella</i> ↑, <i>Clostridium</i> ↑ <i>Lactobacillus</i> ↑, <i>Streptococcus</i> ↑, <i>Pseudomonas</i> ↑, <i>Bacteroidetes</i> ↓, <i>Sutterella</i> ↓ <i>Oscillospira</i> ↓, <i>Faecalibacterium</i> ↓	Tang/2017 ²⁶
PBC (n=39)	Healthy vs PBC	<i>Lactobacillales</i> ↑ <i>Clostridium</i> subcluster XIVa ↓	Abe/2018 ²¹
Oral microbiota			
Disease	Comparison	Genus (or Order)	Author/year
AIH (n=17)	Healthy vs AIH	<i>Veillonella</i> ↑, <i>Streptococcus</i> ↓, <i>Fusobacterium</i> ↓	Abe/2018 ²¹
PBC (n=39)	Healthy vs PBC	<i>Veillonella</i> ↑, <i>Eubacterium</i> ↑ <i>Fusobacterium</i> ↓	Abe/2018 ²¹




Fig. 1. Analysis of the gut and oral microbiota of the PBC, AIH and HC groups based on the T-RFLP profiles. (a) Mean genus abundance of oral microbiota in the PBC, AIH and HC groups. The plotted values are the mean abundance values of the 8 abundant genera in each group. (b) Mean genus or order abundance of gut microbiota in the PBC, AIH and HC groups. The plotted values are the mean abundance values of the 8 abundant genera and 1 abundant order in each group. The results are expressed as the means \pm SDs. Differences were compared using the Mann-Whitney U test; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.0005$.

a decreased abundance of the genus *Clostridium* subcluster XIVa²¹ (Figure 1b). Interestingly, our study suggested that the relative abundance of *Lactobacillales* in feces was positively correlated with the relative abundance of *Veillonella* in saliva from patients with AIH (Figure 2). *Lactobacillus* species had relatively large effect sizes in Behcet's disease microbiota, consistent with the inductive effect of *Lactobacillus* on systemic inflammation. Animal studies using germ-free mice reported that some bacterial species independently promoted arthritis by activating Th17 cells^{22,23}. Dysbiosis of the oral

microbiota is directly and/or indirectly related to the gut microbiota and may be correlated with disease onset.

Gut and oral microbiota in PBC

PBC is a progressive autoimmune liver disease characterized by portal inflammation, immune-mediated destruction of the intrahepatic bile ducts, and the presence of highly specific anti-mitochondrial antibodies in serum^{24,25}. Tang R *et al.* reported that the 16S rRNA gene was analyzed longitudinally by next-generation sequencing (NGS) in PBC patients before treatment with ursodeoxycholic acid (UDCA) using stools of 114 healthy subjects matched in age, gender and body mass index with 79 patients with naive PBC. The number and diversity of observed operational taxonomic units (OTUs) were significantly reduced in the naive PBC patients. The results of principal component analysis by calculating the UniFrac distance showed that the distribution was significantly different between the naive patients with PBC and the healthy subjects. The exploration cohort and the verification cohort had similar results²⁶. Moreover, to investigate the changes in specific gut microbiota components in naive PBC patients, the relative abundance of bacteria at the genus level was compared with that of the healthy subjects. At the genus level, there were changes in the relative abundances of 12 bacteria in naive patients with PBC compared to healthy subjects. Eight bacteria (*Haemophilus*, *Veillonella*, *Clostridium*,




Fig. 2. Correlation between the gut microbiota and oral microbiota in AIH patients.

The relative abundance of *Lactobacillales* in feces was positively correlated with the relative abundance of *Veillonella* in saliva from patients with AIH. The P -value was calculated with the Spearman rank correlation test. $P < 0.05$ was considered significant.

Lactobacillus, *Streptococcus*, *Pseudomonas*, *Klebsiella*, and *Enterobacteriaceae*) had increased relative abundances, and four (*Sutterella*, *Oscillospira*, *Faecalibacterium*, and *Bacteroides*) had decreased relative abundances (Table 1).

Our study indicated that there was a significantly higher frequency of bacteria in the genus *Veillonella* and the genus *Eubacterium* and a lower frequency of bacteria in the genus *Fusobacterium* in the PBC group than in the healthy subjects (Figure 1a). Moreover, the changes in the gut microbiota composition in PBC were characterized by an increase in the abundance of the order *Lactobacillales* and by a decrease in the abundance of the genus *Clostridium* subcluster *XIVa* (Figure 1b). The relative abundance of *Streptococcus* was negatively correlated with the levels of IL-1 β , IL-4, IL-6, IL-7, IL-8, IL-12p70, IL-17, G-CSF, IFN- γ and TNF- α , while the relative abundances of *Veillonella* and *Prevotella/Porphyromonas* were positively correlated with the IgA level in the saliva of patients with PBC. Moreover, the relative abundances of *Neisseria* and *Eubacterium/Filifactor* were positively correlated with the salivary cytokine levels in patients with PBC²¹⁾.

In 37 patients with naive PBC, UDCA was orally administered for 6 months, and changes in the gut microbiota were evaluated before and after treatment. The abundances of *Haemophilus*, *Veillonella*, *Streptococcus*, and *Pseudomonas* decreased after UDCA treatment, and those of *Sutterella*, *Oscillospira*, and *Bacteroides* increased after UDCA treatment in PBC patients²⁶⁾.

Conclusion

These findings suggest that the gut and oral microbiota may play different roles in the pathophysiology of AIH and PBC. Further studies of the establishment and modification of the gut and oral microbiota structure may contribute to the development of a therapeutic strategy for patients with autoimmune liver disease.

Conflict of interest disclosure

The authors declare no conflicts of interest associated with this manuscript.

References

- Rai R, Saraswat VA, Dhiman RK. Gut microbiota: its role in hepatic encephalopathy. *J Clin Exp Hepatol*, **5** : 29-36, 2015.
- Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, *et al.* A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*, **464** : 59-65, 2010.
- Hooper LV, Macpherson AJ. Immune adaptations that maintain homeostasis with the intestinal microbiota. *Nat Rev Immunol*, **10** : 159-169, 2010.
- Fraher MH, O'Toole PW, Quigley EM. Techniques used to characterize the gut microbiota: a guide for the clinician. *Nat Rev Gastroenterol Hepatol*, **9** : 312-22, 2012.
- Avila M, Ojcius DM and Yilmaz, O. The oral microbiota: living with a permanent guest, *DNA Cell Biol*, **28** : 405-411, 2009.
- Dewhirst, FE, Chen, T, Izard, J, Paster BJ, Tanner AC, Yu WH, *et al.* The human oral microbiome. *J. Bacteriol*, **192** : 5002-5017, 2010.
- Curtis MA, Zenobia C. and Darveau RP. The relationship of the oral microbiota to periodontal health and disease. *Cell Host Microbe*, **10** : 302-306, 2011.
- Miyake S, Kim S, Suda W, Oshima K, Nakamura M, Matsuoka T, *et al.* Dysbiosis in the Gut Microbiota of Patients with Multiple Sclerosis, with a Striking Depletion of Species Belonging to Clostridia XIVa and IV Clusters. *PLoS One*, **10**(9) : e0137429, 2015. doi : 10.1371/journal.pone.0137429. PMID : 26367776
- Berer K, Gerdes LA, Cekanaviciute E, Jia X, Xiao L, Xia Z, *et al.* Gut microbiota from multiple sclerosis patients enables spontaneous autoimmune encephalomyelitis in mice. *Proc Natl Acad Sci U S A*, **114**(40) : 10719-10724, 2017. doi : 10.1073/pnas.1711233114. PMID : 28893994
- Cekanaviciute E, Yoo BB, Runia TF, Debelius JW, Singh S, Nelson CA, *et al.* Gut bacteria from multiple sclerosis patients modulate human T cells and exacerbate symptoms in mouse models. *Proc Natl Acad Sci U S A*, **114**(40) : 10713-10718, 2017. doi : 10.1073/pnas.1711235114. PMID : 28893978
- Zhang X, Zhang D, Jia H, Feng Q, Wang D, Liang D, *et al.* The oral and gut microbiomes are perturbed in rheumatoid arthritis and partly normalized after treatment. *Nat Med*, **21**(8) : 895-905, 2015. doi : 10.1038/nm.3914. PMID : 26214836
- Phillips R. Rheumatoid arthritis: Microbiome reflects status of RA and response to therapy. *Nat Rev Rheumatol*, **11**(9) : 502, 2015. doi : 10.1038/nrrheum.2015.109. PMID : 26241185
- Said HS, Suda W, Nakagome S, Chinen H, Oshima K, Kim S, *et al.* Dysbiosis of salivary microbiota in inflammatory bowel disease and its association with oral immunological biomarkers. *DNA Res*, **21**(1) : 15-25, 2014. doi : 10.1093/dnares/dst037. PMID : 24013298

14. Atarashi K, Suda W, Luo C, Kawaguchi T, Motoo I, Narushima S, *et al.* Ectopic colonization of oral bacteria in the intestine drives TH1 cell induction and inflammation. *Science*, **358**(6361): 359-365, 2017. doi: 10.1126/science.aan4526. PMID: 29051379
15. Bajaj JS, Betrapally NS, Hylemon PB, Heuman DM, Daita K, White MB, *et al.* Salivary microbiota reflects changes in gut microbiota in cirrhosis with hepatic encephalopathy. *Hepatology*, **62**(4): 1260-1271, 2015. doi: 10.1002/hep.27819. PMID: 25820757
16. Krawitt EL. Autoimmune hepatitis. *N Engl J Med*, **334**: 897-903, 1996.
17. Lin R, Zhou L, Zhang J, Wang B. Abnormal intestinal permeability and microbiota in patients with autoimmune hepatitis. *Int J Clin Exp Pathol*, **8**(5): 5153-5160, 2015. PMID: 26191211
18. Yuksel M, Wang Y, Tai N, Peng J, Guo J, Beland K, *et al.* A novel "humanized mouse" model for autoimmune hepatitis and the association of gut microbiota with liver inflammation. *Hepatology*, **62**(5): 1536-1550, 2015. doi: 10.1002/hep.27998. PMID: 26185095
19. Czaja AJ. Factoring the intestinal microbiome into the pathogenesis of autoimmune hepatitis. *World J Gastroenterol*, **22**(42): 9257-9278, 2016. Review. PMID: 27895415
20. Wei Y, Li Y, Yan L, Sun C, Miao Q, Wang Q, Xiao X, Lian M, Li B, Chen Y, Zhang J, Li Y, Huang B, Li Y, Cao Q, Fan Z, Chen X, Fang JY, Gershwin ME, Tang R, Ma X. Alterations of gut microbiome in autoimmune hepatitis. *Gut*. 2019. pii: gutjnl-2018-317836. doi: 10.1136/gutjnl-2018-317836. [Epub ahead of print]
21. Abe K, Takahashi A, Fujita M, Imaizumi H, Hayashi M, Okai K, *et al.* Dysbiosis of oral microbiota and its association with salivary immunological biomarkers in autoimmune liver disease. *PLoS One*, **13**(7): e0198757, 2018 Jul 3.
22. Wu HJ, Ivanov II, Darce J, Hattori K, Shima T, Umesaki Y, *et al.* Gut-residing segmented filamentous bacteria drive autoimmune arthritis via T helper 17 cells. *Immunity*, **32**: 815-827, 2010. doi: 10.1016/j.immuni.2010.06.001. PMID: 20620945.
23. Shimizu J, Kubota T, Takada E, Takai K, Fujiwara N, Arimitsu N, *et al.* Bifidobacteria Abundance-Featured Gut Microbiota Compositional Change in Patients with Behcet's Disease. *PLoS One*, **11**: e0153746, 2016. doi: 10.1371/journal.pone.0153746. eCollection 2016. PMID: 27105322
24. Kaplan MM, Gershwin ME. Primary biliary cirrhosis. *N Engl J Med*, **353**: 1261-1273, 2005.
25. Selmi C, Bowlus CL, Gershwin ME, Coppel RL. Primary biliary cirrhosis. *Lancet*, **377**: 1600-1609, 2011.
26. Tang R, Wei Y, Li Y, Chen W, Chen H, Wang Q, Yang F, Miao Q, Xiao X, Zhang H, Lian M, Jiang X, Zhang J, Cao Q, Fan Z, Wu M, Qiu D, Fang JY, Ansari A, Gershwin ME, Ma X. Gut microbial profile is altered in primary biliary cholangitis and partially restored after UDCA therapy. *Gut*, **67**: 534-541, 2018.